

## SÍRIA

*It is a Portuguese variety spread particularly in the region of Alentejo, on the border with Spain, where it is known as Roupeiro. It is also known as Códega in the region of Douro and Alvarurão in Dão.*



**Ampelographic characters:** the bud has an open whitish-green apex with a very intense anthocyanin pigmentation and high density of prostrate hairs. The medium-sized pentagonal leaf is five-lobed, with a slightly open V-shaped petiolar sinus. The cluster is medium-large and compact, with a medium-sized stem. The berry is medium-sized, uniform, short and elliptical.

**Cultural aptitude:** average-high vigorous vine, of semi-upright growth habit, suitable for warm environments.

**Training systems and pruning:** it adapts to different training systems, preferring long pruning (single-cane or double-cane Guyot).

**Bud-burst period:** average-early.

**Ripening period:** late.

**Yield:** average.

**Susceptibility to diseases and adverse conditions:** medium susceptibility to downy mildew, susceptible to powdery mildew, botrytis and mites.

**Enological potential:** it produces wines of light-yellow colour, fresh, slightly acidic and fruity, for preferably young consumption.



### CULTIVATED AREA IN PORTUGAL

YEAR	1989	2000	2011
HECTARES	4,398	11,700	5,040

Vineyards in Alentejo (Portugal)

